

TOPOLOGICAL RIGIDITY OF HIGHER GRAPH MANIFOLDS

NOÉ BÁRCENAS, DANIEL JUAN-PINEDA AND PABLO SUÁREZ-SERRATO

ABSTRACT. In this short note we prove the Borel conjecture for a family of aspherical manifolds that includes higher graph manifolds.

1. INTRODUCTION

The Borel conjecture is a statement about topological rigidity. It states that a homotopy equivalence between two aspherical manifolds is homotopic to a homeomorphism.

A lot of work in geometric topology has been done in the last years with the aim to prove the Borel conjecture using methods involving controlled topology and algebraic K -theory. In particular, the Borel conjecture was shown by R. Frigerio, J.-F. Lafont, and A. Sisto to hold for the class of graph manifolds studied in [11].

On the other hand, relationships between several generalizations of the concept of finite asymptotic dimension in connection with isomorphism conjectures, in algebraic K and L -theory, as well as coarse versions of these have been carried out by G. Carlsson and B. Goldfarb in [7], [12], [8].

The method of proof of the Borel conjecture in this note uses these previous developments. Consider the following construction of smooth n -manifolds M , for $n \geq 3$:

- Definition 1.** (1) For every $i = 1, \dots, r$ take a complete finite-volume non-compact pinched negatively curved n_i -manifold V_i , where $2 \leq n_i \leq n$.
- (2) Denote by M_i the compact smooth manifold with boundary obtained by “truncating the cusps” of V_i , i.e. by removing from V_i a (nonmaximal) horospherical open neighborhood of each cusp.
- (3) Take fiber bundles $Z_i \rightarrow M_i$ with fiber a compact quotient N_i of an aspherical simply connected Lie group \widetilde{N}_i by the action of a uniform lattice Γ_i , of dimension $n - n_i$, i.e. N_i is diffeomorphic to \widetilde{N}_i/Γ_i , where \widetilde{N}_i is a simply connected Lie group and Γ_i is a uniform lattice.
- (4) Fix a complete pairing of diffeomorphic boundary components between distinct Z_i 's, provided one exists, and glue the paired boundary components using diffeomorphisms, to obtain a connected manifold of dimension n .

We will call the Z_i 's the pieces of M and whenever $\dim(M_i) = n$, then we say $Z_i = M_i$ is a pure piece, (short for purely negatively curved).

Remark 1. The construction in the previous definition includes:

- (1) The class of **generalized graph manifolds** of Frigerio, Lafont and Sisto [11]. The pieces V_i in item (1) above are required to be hyperbolic with toral boundary cusps, the N_i in item (3) are required to be tori, and the gluing diffeomorphisms in item (4) are required

to be affine diffeomorphisms. These authors produce examples of manifolds within this class that do not admit any CAT(0) metric.

- (2) The family of **cuspidal decomposable manifolds** of T. Tam Nguyen Phan [14], where interesting (non)rigidity properties are explored. These manifolds only have pure pieces.
- (3) The **affine twisted doubles** of hyperbolic manifolds, for which C.S. Aravinda and T. Farrell study in [1] the existence of nonpositively curved metrics.
- (4) The **higher graph manifolds** studied in [9] by C. Connell and the third named author. In that family, item (3) consists of infanilmanifold bundles with affine structure group, which are moreover trivial near the 'cusp boundary' of the negatively curved pieces in the base. In item (4) the glueing diffeomorphisms are restricted to those which are isotopic to affine diffeomorphisms. These two restrictions are used in [9] to prove statements about collapsing and computations of minimal volume. They turn out not to be needed in the arguments we present for the Borel conjecture to hold true.

The following theorem is our main result:

Theorem 1. *Let M be an n -dimensional manifold constructed as in Definition 1, for $n \geq 6$, then M satisfies the Borel conjecture. That is, given a homotopy equivalence $f : M \rightarrow M'$, where M' is an aspherical n -dimensional manifold, then f is homotopic to a homeomorphism.*

The following section explains the notions of asymptotic dimension, weak regular coherence, and finite decomposition complexity. In the last section a proof of Theorem 1 that uses these properties can be found, and also a proof that presents a slight extension of the general strategy proposed by Frigerio-Lafont-Sisto, and we verify it for the higher graph manifolds whose pieces are trivial bundles.

Acknowledgements. The authors thank Boris Goldfarb and Jim Davis for conversations related to this paper. The first named author acknowledges the support of UNAM PAPIIT Grant ID100315. The second named author has support from UNAM-PAPIIT and CONACyT research grants. The third named author thanks CONACyT Mexico and DGAPA-UNAM for supporting various research initiatives.

2. FINITE ASYMPTOTIC DIMENSION AND WEAK REGULAR COHERENCE

2.1. **Asphericity.** Consider the next definition, following [11], which will be used later on:

Definition 2. *The boundaries of the pieces Z_i that are identified together in Definition 1 will be called the **internal walls** of M .*

Now we will prove, via an adaptation of the arguments of Frigerio-Lafont-Sisto, that the manifolds we are interested in are in fact aspherical.

Lemma 2. *If M is a manifold (possibly with boundary) constructed as in Definition 1, then M is aspherical.*

Proof. This proof is by induction on the number of internal walls c of M . If $c = 0$ then $M = Z$ for some bundle Z over a closed, negatively curved base. It follows from the homotopy exact sequence for the bundle Z that M is aspherical in this case, establishing the base case for our inductive argument.

Assume $c > 0$, and that the result holds for manifolds constructed as in Definition 1, with strictly less than c internal walls. Cut open M along an arbitrary internal wall W . Our inductive hypothesis implies that now M is obtained by gluing one or two (depending on whether W separates M or not) aspherical spaces. Since the inclusion of W in the piece(s) in M it belongs to is π_1 -injective, it follows from a classical result of Whitehead [16] that M is aspherical. \square

2.2. Finite asymptotic dimension. Let G be a finitely presented group. Fix a finite generator set S and consider the word metric d_S induced by the generating set. With this metric, the group G is a proper metric space.

Definition 3. A family $\{U\}$ of subsets in a metric space X is D -disjoint if $d(U, U') > D$ for all subsets in the family. The asymptotic dimension $\text{asdim} X$ of X is the smallest number n such that for any $D > 0$ there is a uniformly bounded cover of X by $n + 1$ -families of D -disjoint families of subsets.

An example of spaces (and groups) for which their asymptotic dimension can be explicitly computed are precisely quotients of simply connected Lie groups:

Theorem 3. (Carlsson-Goldfarb, Cor. 3.6 in [6]) Let Γ be a cocompact lattice in a connected Lie group G with maximal compact subgroup K . Then $\text{asdim} \Gamma = \dim(G/K)$.

For spaces that are built up using smaller subsets, there is a theorem that allows us to bound the asymptotic dimension of the total space. Let X be a metric space. The family $\{X_\alpha\}$ of subsets of X is said to satisfy the inequality $\text{asdim} X_\alpha \leq n$ **uniformly** if for every $r < \infty$ a constant R can be found so that for every α there exists R -disjoint families $U_\alpha^0, U_\alpha^1, U_\alpha^2, \dots, U_\alpha^n$ of R -bounded subsets of X_α covering X_α .

Theorem 4. (Union theorem, Bell-Dranishnikov, Thm. 25 in [3]) Let $X = \bigcup_\alpha X_\alpha$ be a metric space where the family $\{X_\alpha\}$ satisfies the inequality $\text{asdim} X_\alpha \leq n$ uniformly. Suppose further that for every r there is a $Y_r \subset X$ with $\text{asdim} Y_r \leq n$ so that $d(X_\alpha - Y_r, X_{\alpha'} - Y_r) \geq r$ whenever $X_\alpha \neq X_{\alpha'}$. Then $\text{asdim} X \leq n$.

Lemma 5. The fundamental group $\pi_1(M)$ of a manifold M of dimension n constructed as in Definition 1 has finite asymptotic dimension.

Proof. The fundamental groups of the pieces $\pi_1(Z_i)$ fit in an exact sequence

$$1 \rightarrow \pi_1(N_i) \rightarrow \pi_1(Z_i) \rightarrow \pi_1(M_i) \rightarrow 1.$$

The asymptotic dimension of $\pi_1(M_i)$ equals $n - n_i$ by the Cartan-Hadamard Theorem. On the other hand, the asymptotic dimension of the fibres, which are quotients of Lie groups G under the action of a uniform lattice, equals $\dim(G/K) < \infty$ by Theorem 3.

Finally, we invoke Theorem 4, from which we conclude that the asymptotic dimension of $\pi_1(X)$ is finite. \square

2.3. Finite decomposition complexity. We will briefly define the notion of *straight finite decomposition complexity*, since we use it as a key property in the proof of the main result presented below.

Let \mathcal{X} and \mathcal{Y} be two families of metric spaces, and $R > 0$. The family \mathcal{X} is called R -decomposable over \mathcal{Y} if, for any space X in \mathcal{X} there are collections of subsets $\{U_{1,\alpha}\}$ and $\{U_{2,\beta}\}$ such that

$$X = \bigcup_{i=1,2,\gamma=\alpha,\beta} U_{i,\gamma}.$$

Each $U_{i,\gamma}$ is a member of the family \mathcal{Y} , and each of the collections $\{U_{1,\alpha}\}$ and $\{U_{2,\beta}\}$ is R -disjoint. A family of metric spaces is called *bounded* if there is a uniform bound on the diameters of the spaces in the family.

Definition 4. *A metric space X has straight finite decomposition complexity if, for any sequence $R_1 \leq R_2 \leq \dots$ of positive numbers, there exists a finite sequence of metric families V_1, V_2, \dots, V_n such that X is R_1 -decomposable over V_1 , X is R_2 -decomposable over V_2 , etc, and the family V_n is bounded.*

The following well known lemma ties this notion with that of asymptotic dimension, for completeness we include a proof:

Lemma 6. *If a group has finite asymptotic dimension, then it has straight finite decomposition complexity.*

Proof. It was shown by Guentner-Tessera-Yu that a countable metric space of finite asymptotic dimension has finite decomposition complexity in [13]. As part of their study of straight finite decomposition complexity, Dranishnikov-Zarichnyi showed in [10] that groups with finite decomposition complexity have straight finite decomposition complexity. \square

3. TWO PROOFS

3.1. The Carlsson-Goldfarb approach to the Borel conjecture. Let Γ be the fundamental group of a manifold constructed as in Definition 1. The strategy for proving Theorem 1 for manifolds with fundamental group Γ consists of showing that Γ satisfies the following properties:

- (1) Γ has finite asymptotic dimension.
- (2) Γ has a finite model for the classifying space $B\Gamma$.

A group satisfying these two conditions has been proven to also satisfy the integral isomorphism conjecture in algebraic K-theory, according to Theorem 3.11 of Goldfarb in [12].

Proof. (of Theorem 1)

Item (1) was shown in Lemma 5 above.

Item (2) follows from the fact that these are fundamental groups of compact aspherical manifolds (possibly with boundary) Therefore the Borel conjecture holds for the manifolds in Definition 1. \square

This simple strategy provides an alternative to the one layed out by Frigerio-Lafont-Sisto in [11]. We also present in the following a modified version of their strategy, and verify that it can be carried out for certain manifolds within those of Definition 1.

In a series of articles, B. Goldfarb and G. Carlsson have investigated several notions which generalize that of regular coherence for the group ring of infinite groups. The main geometric interest on this situation resides on the fact that these conditions are strong enough to allow the vanishing of the Whitehead group and negative algebraic K-theory groups of group

rings, and weak enough to be handled with methods dealing with coarse versions of the isomorphism conjecture in algebraic K -theory [12],[6].

We will recall some definitions and fundamental results related to finite asymptotic dimension and the coarse assembly map in the boundedly controlled setting. See, for example, [5] and [12] for further reference.

Let $\mathcal{P}(G)$ be the power set viewed as a category where morphisms are inclusions of subsets. Let R be a noetherian ring and consider a finitely generated $R[G]$ -module \mathcal{M} . A G -filtration of \mathcal{M} is a functor $f : \mathcal{P}(G) \rightarrow R\text{-Sub}(\mathcal{M})$ to the category of R -submodules of \mathcal{M} such that $f(G) = \mathcal{M}$, and each bounded set in the word metric d_S , $T \subset G$ is mapped to a finitely generated R -submodule. Such a functor f is equivariant if $f(gS) = gf(S)$

Definition 5. A homomorphism $\phi : F_1 \rightarrow F_2$ between finitely generated $R[G]$ -modules with fixed filtrations f_1, f_2 is boundedly controlled with respect to the bound $D > 0$ if $\phi(f_1(S)) \subset f_2(B_D(S))$ for each subset $S \subset G$. If ϕ also satisfies $\phi F_1 \cap f_2(S) \subset \phi F_1(B_D(S))$, then F is called boundedly bicontrolled.

Definition 6. Let \mathcal{M} be a finitely presented $R[G]$ -module. A finite presentation $F : R[G]^m \rightarrow R[G]^n \rightarrow \mathcal{M}$ is admissible if the homomorphism F is boundedly bicontrolled.

Definition 7. A group ring $R[G]$ is weakly coherent if every $R[G]$ -module with an admissible presentation has a projective resolution of finite type. Similarly, a group ring is weakly regular coherent if every $R[G]$ -module with an admissible presentation has finite homological dimension.

Theorem 7. (Carlsson-Goldfarb, Corollary 3.9 in [12]) Let R be a noetherian ring and let G be a group of finite asymptotic dimension. Then, the group ring $R[G]$ is weakly regular coherent.

Weak regular coherence has been verified to be enough to guarantee the vanishing of Whitehead groups and negative algebraic K -theory.

Theorem 8. (Goldfarb, Theorem 3.11 in [12]). Let G be a group of finite asymptotic dimension (or more generally of finite decomposition complexity, as explained in [12]). Assume that there is a finite model for the classifying space $K(G, 1)$. Then, the assembly map in algebraic K -theory is an isomorphism. In particular, the Whitehead group of G vanishes.

As a consequence of theorem 8 and lemma 5, we obtain:

Corollary 9. The group ring $\mathbb{Z}\pi_1(M)$ of a manifold M constructed as in Definition 1 is weakly regular coherent.

3.2. An extension of the Frigerio-Lafont-Sisto approach to the Borel conjecture.

The proof of the Borel conjecture for the class of manifolds studied by Frigerio-Lafont-Sisto in [11] in fact developed a general strategy to be carried out for a given family of manifolds. In their Theorem 3.1 they proved that if a manifold is built up from a geometric decomposition, as are the higher graph manifolds in this paper, and satisfies the following six conditions, then it also satisfies the Borel conjecture:

- (1) Each of the inclusions $W_{i,j} \rightarrow Z_i$ is π_1 -injective.
- (2) Each of the pieces Z_i and each of the walls $W_{i,j}$ are aspherical.
- (3) Each of the pieces Z_i and each of the walls $W_{i,j}$ satisfy the Borel Conjecture.

- (4) The rings $\mathbf{Z}\pi_1(W_{i,j})$ are all regular coherent.
- (5) $Wh_k(\mathbf{Z}\pi_1(W_{i,j})) = 0 = Wh_k(\mathbf{Z}\pi_1(Z_i))$ for $k \leq 1$.
- (6) Each of the inclusions $\pi_1(W_{i,j}) \rightarrow \pi_1(Z_i)$ is square-root-closed.

We propose a slightly modified version of this strategy, where we replace the last three conditions, (4), (5) and (6), by a couple of new requirements. So that we obtain the following:

Lemma 10. *Let M be a compact manifold of dimension $n \geq 6$ with a topological decomposition (as described in [11]). Assume the following conditions hold:*

- (1) *Each of the inclusions $W_{i,j} \rightarrow Z_i$ is π_1 -injective.*
- (2) *Each of the pieces Z_i and each of the walls $W_{i,j}$ are aspherical.*
- (3) *Each of the pieces Z_i and each of the walls $W_{i,j}$ satisfy the Borel Conjecture.*
- (4) *The group $\Gamma = \pi_1(M)$ has finite decomposition complexity.*
- (5) *There exists a finite model for the classifying space $K(\Gamma, 1)$.*

Then the manifold M also satisfies the Borel conjecture.

Proof. Conditions (4) and (5) imply that the Whitehead groups $Wh_i(\mathbf{Z}\Gamma) = 0$, for $i \leq 1$, as proved in [12].

Therefore the rest of the proof presented in Theorem 3.1 [11] follows through, and the result holds. \square

Now we will concentrate on certain higher graph manifolds, explained briefly in the introduction (see [9]).

Lemma 11. *Assume M is a higher graph manifold, all of whose pieces are trivial as bundles. Then, each of the pieces $Z_i \cong N_i \times M_i$, and each of the walls $W_{i,j}$, satisfy the fibred isomorphism conjecture (FIC) of Farrell-Jones.*

Proof. First notice that the validity of FIC for the walls $W_{i,j}$ follows from the work of Bartels-Farrell-Lück in [2], since these are quotients of Lie groups (see also their Remark 2.13).

As each piece Z_i is a trivial fibre bundle

$$Z_i \cong N_i \times M_i$$

the fundamental group of Z_i is a product

$$\pi_1(Z_i) \cong \pi_1(N_i) \times \pi_1(M_i).$$

Recall that M_i is a manifold that admits a pinched negatively curved metric. So it also admits a $CAT(0)$ metric, and therefore FIC holds for $\pi_1(M_i)$. The fibres satisfy FIC following [2]. Therefore $\pi_1(Z_i)$ also satisfies FIC, by Theorem 2.9 in [2]. \square

As a consequence we obtain that the Borel conjecture holds for each of the pieces Z_i , with trivial fibration structure, and each of the walls $W_{i,j}$, and so condition (3) is verified.

Lemma 12. *Let M be a higher graph manifold, all of whose pieces Z_i are trivial as bundles, and let $W_{i,j}$ denote its internal walls. Then, the rings $\mathbf{Z}\pi_1(W_{i,j})$ are weakly regular coherent.*

Proof. From the proof of Lemma 5, we conclude that these groups have finite asymptotic dimension. Now the result follows from Theorem 7. \square

Lemma 13. *Let M be a higher graph manifold, all of whose pieces Z_i are trivial as bundles, and let $W_{i,j}$ denote its internal walls. Then, $Wh_k(\mathbf{Z}\pi_1(W_{i,j})) = 0 = Wh_k(\mathbf{Z}\pi_1(Z_i))$ for $k \leq 1$.*

Proof. Since each of the walls and pieces are aspherical, their fundamental groups are torsion-free. By the previous Lemma 11, the result holds for each of the pieces and walls. Alternatively, the result follows from Theorem 8. \square

Lemma 14. *Let M be a manifold constructed as in Definition 1. For every $1 \leq i \leq r$, the map $N_i \rightarrow X$ is π_1 -injective. Moreover, the image of $\pi_1(N_i)$ is a square root closed subgroup in the group $\pi_1(X)$.*

Proof. Consider the long exact sequence of homotopy groups of a fibration:

$$\dots \rightarrow \pi_n(N_i) \rightarrow \pi_n(Z_i) \rightarrow \pi_n(M_i) \rightarrow \dots$$

The connectedness of N_i implies the π_1 -injectivity condition.

Using proposition VII.2 in page 168 of [4], it suffices to verify the square root closed condition in the fundamental groups of the edges $\pi_1(W_i, j) \rightarrow \pi_1(Z_i)$. Using the long exact sequence of the fibration again, this is equivalent to showing that there are no 2-torsion elements in $\pi_1(M_i)$. This is certainly the case, since M_i is an aspherical manifold. \square

Lemma 15. *Let M be a manifold constructed as in Definition 1 and $\Gamma = \pi_1(M)$. Then there exists a finite model for $K(\Gamma, 1)$.*

Proof. Notice that the manifold M is aspherical and hence it is itself a finite model for $K(\Gamma, 1)$. \square

Now we collect all of these auxiliary results to present:

Theorem 16. *Let M be a higher graph manifold of dimension ≥ 6 . Assume that all of the pieces of M are trivial bundles. Then M satisfies the Borel conjecture.*

Proof. Notice that these higher graph manifolds satisfy all the hypothesis of Lemma 10. As has been shown in Lemmas 6, 11, 13, 14, and 15. \square

REFERENCES

- [1] C.S. Aravinda, T. Farrell, *Twisted doubles and nonpositive curvature*, Bull. London Math. Soc. 41 (2009) 1053–1059.
- [2] A. Bartels, F.T. Farrell, W. Lück, *The Farrell-Jones conjecture for cocompact lattices in virtually connected Lie groups*, Journal of the A.M.S., 27 (2014), no. 2, 339–388.
- [3] G. Bell, A. Dranishnikov, *Asymptotic dimension*, Topology Appl. 155 (2008), no. 12, 1265–1296.
- [4] S. Capell, *Splitting theorems for manifolds* Invent. Math. 33 (2) 1976, 69–170.
- [5] G. Carlsson, B. Goldfarb, *The integral Novikov Conjecture for groups with finite asymptotic dimension*, Invent. Math. 157 (2004), 405–418.
- [6] G. Carlsson, B. Goldfarb, *On homological coherence of discrete groups*, J. of Algebra 276 (2) 2004 502–514.
- [7] G. Carlsson, B. Goldfarb, *Algebraic K-Theory of Geometric Groups*, arXiv:1305.3349.
- [8] G. Carlsson, B. Goldfarb, *Equivariant Stable Homotopy Methods in the Algebraic K-Theory of Infinite Groups*, arXiv:1412.3483.
- [9] C. Connell, P. Suárez-Serrato, *On higher graph manifolds*, arXiv:1208.4876.
- [10] A. Dranishnikov, M. Zarichnyi, *Asymptotic dimension, decomposition complexity, and Haver’s property C*, Topology Appl. 169 (2014), 99–107.
- [11] R. Frigerio, J.-F. Lafont, A. Sisto, *Rigidity of high dimensional graph manifolds*, Astérisque, to appear.
- [12] B. Goldfarb, *Weak coherence and the K-Theory of groups with finite decomposition complexity*, IMRN, to appear, arXiv:1307.5345.
- [13] E. Guentner, R. Tessera, G. Yu, *Discrete groups with finite decomposition complexity*. Groups Geom. Dyn. 7 (2013), no. 2, 377–402.

- [14] T. Tam Nguyen Phan, *Smooth (non)rigidity of cusp-decomposable manifolds*, Comment. Math. Helv. 87 (2012), no. 4, 780–804.
- [15] F. Waldhausen, *Algebraic K-Theory of generalized free products I, II, III, IV*, Ann. of Math (2), 108 (1978) 135–256.
- [16] J.H.C. Whitehead, *On the asphericity of regions in a 3-sphere*, Fund. Math.32 (1939), 149–166.

CENTRO DE CIENCIAS MATEMÁTICAS. UNIVERSIDAD NACIONAL AUTÓNOMA DE MÉXICO, AP.POSTAL 61-3 XANGARI. 58089 MORELIA, MICHOACÁN MÉXICO
E-mail address: `barcenas@matmor.unam.mx`
URL: `http://www.matmor.unam.mx/~barcenas`
E-mail address: `daniel@matmor.unam.mx`

INSTITUTO DE MATEMÁTICAS, UNIVERSIDAD NACIONAL AUTÓNOMA DE MÉXICO, CIUDAD UNIVERSITARIA, COYOACÁN, 04510. MEXICO CITY, D. F. MÉXICO
URL: `http://www.matem.unam.mx/PabloSuarezSerrato`
E-mail address: `pablo@im.unam.mx`