

MATHIAS-PRIKRY AND LAVER-PRIKRY TYPE FORCING

MICHAEL HRUŠÁK AND HIROAKI MINAMI

ABSTRACT. We study the Mathias-Prikry and Laver-Prikry forcings associated with filters on ω . We give a combinatorial characterization of Martin's number for these forcing notions and present a general scheme for analyzing preservation properties for them. In particular, we give a combinatorial characterization of those filters for which the Mathias-Prikry forcing does not add any dominating reals.

INTRODUCTION

In recent years, a variety of consistency results have been given using the Mathias-Prikry and the Laver-Prikry forcing associated with filters.

Let \mathcal{F} be a filter on ω . The *Mathias-Prikry* forcing associated with \mathcal{F} , denoted by $\mathbb{M}_{\mathcal{F}}$ consists of pairs $\langle s, A \rangle$ such that $s \in [\omega]^{<\omega}$, $A \in \mathcal{F}$ and $s \cap A = \emptyset$. The ordering $\langle s, A \rangle \leq \langle t, B \rangle$ if $s \supset t$, $A \subset B$ and $s \setminus t \subset B$.

We will refer to the union of the first coordinates of conditions in the generic filter as the generic subset of ω , and denote it by \dot{a}_{gen} .

The *Laver-Prikry* forcing associated with \mathcal{F} , denoted by $\mathbb{L}_{\mathcal{F}}$ consists of subtrees $T \subset \omega^{<\omega}$ which have a stem $s \in T$ (denoted by $\text{stem}(T)$) such that for every $t \in T$ either $t \subset s$ or $s \subset t$ and for every $t \in T$ extending s the set

$$\text{Succ}_T(t) = \{n \in \omega : t \frown \langle n \rangle \in T\} \in \mathcal{F}.$$

The order on $\mathbb{L}_{\mathcal{F}}$ is given by inclusion.

These forcing notions play a significant role in the use of the matrix iteration introduced by Blass and Shelah [5] and further developed and used by Shelah [20], Brendle [10] and Brendle and Fischer [11].

The Laver-Prikry forcing was used to separate variants of the groupwise density number and the distributivity numbers by Brendle in [7, 8, 9] and by Brendle and Hrušák to show it is relatively consistent that every countable FU_{fin} space of weight \aleph_1 is metrizable [12].

The research of first and second authors was partially supported by PAPIIT grant IN101608 and CONACYT grant 46337.

In [19], Raghavan constructs a model of ZFC without strongly separable almost disjoint families by using a similar technique.

In this paper, we shall study the relation between combinatorial properties of an ideal \mathcal{I} and the forcing properties of the Mathias-Prikry and the Laver-Prikry type forcings associated with the dual-filter \mathcal{I}^* (denoted by $\mathbb{M}_{\mathcal{I}^*}$ and $\mathbb{L}_{\mathcal{I}^*}$ respectively) often expressed in terms of the Katětov order, paying special attention to definable (Borel, analytic) ideals.

Both forcing notions are clearly c.c.c, in fact, σ -centered. Also $\mathbb{L}_{\mathcal{I}^*}$ adds a dominating real (the generic function \dot{f}_{gen} is dominating).

In section 1, we give a combinatorial characterization of the Martin number of $\mathbb{M}_{\mathcal{I}^*}$ and $\mathbb{L}_{\mathcal{I}^*}$ and introduce the separating number of the corresponding ideal. In section 2, we investigate the relationship between preservation statements for $\mathbb{M}_{\mathcal{I}^*}$ and $\mathbb{L}_{\mathcal{I}^*}$ and combinatorial properties of \mathcal{I} . Finally, in section 3, we give a characterization of those ideals \mathcal{I} such that $\mathbb{M}_{\mathcal{I}^*}$ does not add any dominating reals.

For a set X , we call $\mathcal{I} \subset \mathcal{P}(X)$ an *ideal on X* if

- (1) for $A, B \in \mathcal{I}$, $A \cup B \in \mathcal{I}$,
- (2) for $A, B \subset X$, $A \subset B$ and $B \in \mathcal{I}$ implies $A \in \mathcal{I}$ and
- (3) $X \notin \mathcal{I}$.

We assume that all ideals on X contain $[X]^{<\omega}$, all finite subsets of X . If \mathcal{I} is an ideal on X , \mathcal{I}^* is the dual filter, consisting of complements of the sets in \mathcal{I} . \mathcal{I}^+ denotes the collection of \mathcal{I} -positive set, i.e., subsets of X which are not in \mathcal{I} . We say that an ideal \mathcal{I} on the set of all natural numbers ω is *tall* if for each $A \in [\omega]^\omega$ there is a $I \in \mathcal{I}$ such that $I \cap A$ is infinite. If \mathcal{I} is an ideal on ω and $Y \in \mathcal{I}^+$, we denote by $\mathcal{I} \upharpoonright Y$ the ideal $\{I \cap Y : I \in \mathcal{I}\}$ on Y .

The topology of $\mathcal{P}(\omega)$ is induced by identifying each subset of ω with its characteristic function, where 2^ω is equipped with the product topology. We call an ideal \mathcal{I} on ω a *Borel ideal* if \mathcal{I} is Borel in this topology.

Given a tall ideal \mathcal{I} on ω and a forcing notion \mathbb{P} , we say that the forcing \mathbb{P} *destroys* \mathcal{I} if there is a \mathbb{P} -name \dot{x} for an element of $[\omega]^\omega$ such that

$$\Vdash_{\mathbb{P}} \forall I \in \mathcal{I} \cap V (|I \cap \dot{x}| < \aleph_0).$$

We say that a family $\mathcal{K} \subset [\omega]^\omega$ is *countably tall* (or *ω -hitting*) if given $(A_n : n \in \omega) \subset [\omega]^\omega$ there is an $K \in \mathcal{K}$ such that for $n \in \omega$, $|K \cap A_n| = \aleph_0$.

The Katětov order on ideals is defined as follows: Suppose \mathcal{I} and \mathcal{J} are ideals on countable sets X and Y respectively. Then $\mathcal{I} \leq_K \mathcal{J}$ if there exists a function $f : Y \rightarrow X$ such that for each $I \in \mathcal{I}$, $f^{-1}[I] \in \mathcal{J}$.

When dealing with ideals on countable sets, we often use the following cardinal invariants [15]:

$$\begin{aligned} \text{add}^*(\mathcal{I}) &= \min\{|\mathcal{A}| : \mathcal{A} \subset \mathcal{I} \wedge (\forall X \in \mathcal{I})(\exists A \in \mathcal{A})(A \not\subset^* X)\}. \\ \text{cov}^*(\mathcal{I}) &= \min\{|\mathcal{A}| : \mathcal{A} \subset \mathcal{I} \wedge (\forall X \in [\omega]^\omega)(\exists A \in \mathcal{A})(|A \cap X| = \aleph_0)\}. \\ \text{non}^*(\mathcal{I}) &= \min\{|\mathcal{X}| : \mathcal{X} \subset [\omega]^\omega \wedge (\forall I \in \mathcal{I})(\exists X \in \mathcal{X})(|I \cap X| < \aleph_0)\}. \\ \text{cof}^*(\mathcal{I}) &= \min\{|\mathcal{A}| : \mathcal{A} \subset \mathcal{I} \wedge (\forall I \in \mathcal{I})(\exists A \in \mathcal{A})(I \subset^* A)\}^1. \end{aligned}$$

1. MARTIN NUMBERS OF $\mathbb{M}_{\mathcal{I}^*}$ AND $\mathbb{L}_{\mathcal{I}^*}$

Recall that the *Martin number* $\mathfrak{m}(\mathbb{P})$ of a partial order \mathbb{P} is the minimal size of a family of dense open subsets of \mathbb{P} such that no filter on \mathbb{P} intersects with them all.

In this section, we shall give a combinatorial characterization of the cardinal invariants $\mathfrak{m}(\mathbb{L}_{\mathcal{I}^*})$ and $\mathfrak{m}(\mathbb{M}_{\mathcal{I}^*})$.

Both forcings $\mathbb{M}_{\mathcal{I}^*}$, and $\mathbb{L}_{\mathcal{I}^*}$ destroy the ideal \mathcal{I} . In fact, they do more than that. $\mathbb{M}_{\mathcal{I}^*}$ and $\mathbb{L}_{\mathcal{I}^*}$ *separate* \mathcal{I} and \mathcal{I}^+ , that is, they add a set $a_{gen} \subset \omega$ which is almost disjoint from every $I \in \mathcal{I}$, and have infinite intersection with every $X \in \mathcal{I}^+ \cap V$. It is useful to introduce the corresponding cardinal invariant, the separating number of an ideal \mathcal{I} .

Let \mathcal{I} be an ideal on ω . Let $\mathcal{G} \subset \mathcal{I}$, $\mathcal{H} \subset \mathcal{I}^+$ and $A \subset \omega$. We say A *separates* \mathcal{G} from \mathcal{H} if

- (1) $|A \cap I| < \aleph_0$ for $I \in \mathcal{G}$ and
- (2) $|A \cap X| = \aleph_0$ for $X \in \mathcal{H}$.

For an ideal \mathcal{I} , the *separating number* $\text{sep}(\mathcal{I})$ is

$$\begin{aligned} \text{sep}(\mathcal{I}) &= \min\{|\mathcal{G}| + |\mathcal{H}| : \mathcal{G} \subset \mathcal{I} \wedge \mathcal{H} \subset \mathcal{I}^+ \wedge \\ &\quad \forall A \subset \omega \exists I \in \mathcal{G} \exists X \in \mathcal{H} (|A \cap I| = \omega \text{ or } |A \cap X| < \omega)\}. \end{aligned}$$

It is clear from the definition that $\text{add}^*(\mathcal{I}) \leq \text{sep}(\mathcal{I}) \leq \text{cov}^*(\mathcal{I})$ for every tall ideal \mathcal{I} and that $\text{sep}(\mathcal{I}) = \text{cov}^*(\mathcal{I})$ if \mathcal{I} is a maximal ideal.

Proposition 1.1. [15] *Let \mathcal{I} and \mathcal{J} be ideals on ω . Suppose \mathcal{I} is below \mathcal{J} in the Rudin-Kiesler order, that is, there exists $f : \omega \rightarrow \omega$ such that for every $A \subset \omega$, $A \in \mathcal{I}$ if and only if $f^{-1}[A] \in \mathcal{J}$. Then $\text{sep}(\mathcal{I}) \leq \text{sep}(\mathcal{J})$.*

¹In [13], Brendle and Shelah introduced cardinal invariants $\mathfrak{p}(\mathcal{F})$ and $\pi\mathfrak{p}(\mathcal{F})$ associated with an ultrafilter \mathcal{F} . For all tall ideals \mathcal{I} , $\text{add}^*(\mathcal{I}) = \mathfrak{p}(\mathcal{I}^*)$, $\text{cov}^*(\mathcal{I}) = \pi\mathfrak{p}(\mathcal{I}^*)$, $\text{non}^*(\mathcal{I}) = \pi_\chi(\mathcal{I}^*)$ and $\text{cof}^*(\mathcal{I}) = \chi(\mathcal{I}^*)$.

Brendle and Shelah characterized the Martin number of the Mathias-Prikry and Laver-Prikry type for ultrafilters in [13].

Theorem 1.2. [13] *Let \mathcal{U} be an ultrafilter. Then*

- (1) $\mathfrak{m}(\mathbb{M}_{\mathcal{U}}) = \text{cov}^*(\mathcal{U}^*)$.
- (2) $\mathfrak{m}(\mathbb{L}_{\mathcal{U}}) = \min\{\mathfrak{b}, \text{cov}^*(\mathcal{U}^*)\}^2$.

We will prove analogous results for arbitrary filter/ideal.

1.1. Martin number of $\mathbb{L}_{\mathcal{I}^*}$. Recall that an ultrafilter \mathcal{U} on ω is *nowhere dense* if for every function $f : \omega \rightarrow \mathbb{R}$ there is a $U \in \mathcal{U}$ such that $f[U]$ is a nowhere dense subset of \mathbb{R} . It is known (see [3]) that the Lave-Prikry forcing with \mathcal{U} adds a Cohen real if and only if \mathcal{U} is not a nowhere dense ultrafilter.

The following was announced in [15].

Theorem 1.3. *For every ideal \mathcal{I} on ω ,*

$$\mathfrak{m}(\mathbb{L}_{\mathcal{I}^*}) = \begin{cases} \min\{\mathfrak{b}, \text{sep}(\mathcal{I})\} & \text{if } \mathcal{I}^* \text{ is a nowhere dense ultrafilter,} \\ \min\{\text{add}(\mathcal{M}), \text{sep}(\mathcal{I})\} & \text{otherwise.} \end{cases}$$

Proof. (i) If \mathcal{I}^* is a nowhere dense ultrafilter, then the required statement holds by Theorem 1.2 (2) as $\text{sep}(\mathcal{I}^*) = \text{cov}^*(\mathcal{U}^*)$ for every ultrafilter \mathcal{U} .

Suppose that \mathcal{I}^* is not a nowhere dense ultrafilter. First we shall show $\mathfrak{m}(\mathbb{L}_{\mathcal{I}^*}) \leq \min\{\text{add}(\mathcal{M}), \text{sep}(\mathcal{I})\}$.

Since $\mathbb{L}_{\mathcal{I}^*}$ adds a dominating real, $\mathfrak{m}(\mathbb{L}_{\mathcal{I}^*}) \leq \mathfrak{b}$. Since \mathcal{I}^* is not a nowhere dense ultrafilter, $\mathbb{L}_{\mathcal{I}^*}$ adds a Cohen real (see [3]). So $\mathfrak{m}(\mathbb{L}_{\mathcal{I}^*}) \leq \text{cov}(\mathcal{M})$. Since $\text{add}(\mathcal{M}) = \min\{\mathfrak{b}, \text{cov}(\mathcal{M})\}$, $\mathfrak{m}(\mathbb{L}_{\mathcal{I}^*}) \leq \text{add}(\mathcal{M})$.

To see that $\mathfrak{m}(\mathbb{L}_{\mathcal{I}^*}) \leq \text{sep}(\mathcal{I})$, suppose that $\kappa < \mathfrak{m}(\mathbb{L}_{\mathcal{I}^*})$ and let $\mathcal{J} \subset \mathcal{I}$ and $\mathcal{H} \subset \mathcal{I}^+$ such that $|\mathcal{J}| + |\mathcal{H}| \leq \kappa$. For $J \in \mathcal{J}$, put

$$D_J = \{T \in \mathbb{L}_{\mathcal{I}^*} : \forall t \in T(\text{stem}(T) \subset t \rightarrow \text{Succ}_T(t) \cap J = \emptyset)\}.$$

For $H \in \mathcal{H}$ and $n \in \omega$, define

$$E_{H,n} = \{T \in \mathbb{L}_{\mathcal{I}^*} : \text{rang}(\text{stem}(T)) \cap H \setminus n \neq \emptyset\}.$$

Then D_J and $E_{H,n}$ are dense for $J \in \mathcal{J}$, $H \in \mathcal{H}$ and $n \in \omega$.

Let $G \subset \mathbb{L}_{\mathcal{I}^*}$ be a $\{D_J : J \in \mathcal{J}\} \cup \{E_{H,n} : H \in \mathcal{H} \wedge n \in \omega\}$ -generic. Let $f_G = \bigcup \{\text{stem}(T) : T \in G\}$. By genericity, $\text{rang}(f_G) \cap J$ is finite for $J \in \mathcal{J}$ and $\text{rang}(f_G) \cap H$ is infinite for $H \in \mathcal{H}$. So $\text{rang}(f_G)$ separates \mathcal{J} from \mathcal{H} . Therefore $\kappa < \text{sep}(\mathcal{I})$.

(ii) $\min\{\text{add}(\mathcal{M}), \text{sep}(\mathcal{I})\} \leq \mathfrak{m}(\mathbb{L}_{\mathcal{I}^*})$.

⁰Brendle and Shelah investigated cardinal invariants of ideals $\ell_{\mathcal{U}}^0$ and $r_{\mathcal{U}}^0$ associated with an ultrafilter \mathcal{U} . For all ideals \mathcal{I} , $\text{cov}(\ell_{\mathcal{I}^*}^0) = \mathfrak{m}(\mathbb{L}_{\mathcal{I}^*})$ and $\text{cov}(r_{\mathcal{I}^*}^0) = \mathfrak{m}(\mathbb{M}_{\mathcal{I}^*})$.

Suppose $\kappa < \text{add}(\mathcal{M}), \text{sep}(\mathcal{I})$. Let $\{D_\alpha : \alpha < \kappa\}$ be a family of dense open subsets of $\mathbb{L}_{\mathcal{I}^*}$.

For each $\alpha < \kappa$, let $\{T_\alpha^n : n \in \omega\}$ be a maximal antichain of D_α . Let $I_{\alpha,t}^n = \omega \setminus \text{Succ}_{T_\alpha^n}(t) \in \mathcal{I}$ for $\alpha < \kappa, n \in \omega$ and $t \in T_\alpha^n$ with $t \supset \text{stem}(T_\alpha^n)$.

Fix $\alpha < \kappa$. Define a rank function $\text{rk}_\alpha : \omega^{<\omega} \rightarrow \text{Ord}$ by

- (1) $\text{rk}_\alpha(t) = 0$ if $\exists n \in \omega (t \in T_\alpha^n \text{ and } t \supset \text{stem}(T_\alpha^n))$.
- (2) $\text{rk}_\alpha(t) \leq \beta$ if $H_t^\alpha = \{n \in \omega : \text{rk}_\alpha(t \frown \langle n \rangle) < \beta\} \in \mathcal{I}^+$.

Claim 1.4. [12, Lemma 4] *For all $t \in \omega^{<\omega}$, $\text{rk}_\alpha(t)$ is defined.*

Since $\kappa < \text{sep}(\mathcal{I})$, there is an $A \in [\omega]^\omega$ such that for every $\alpha < \kappa, n \in \omega, t \in T_\alpha^n$ and $s \in \omega^{<\omega}$, $|A \cap I_{\alpha,t}^n| < \aleph_0$, i.e., $A \subset^* \text{Succ}_{T_\alpha^n}(t)$ and $|A \cap H_s^\alpha| = \aleph_0$.

Let $\mathbb{L}_{\text{fin}(A)^*}$ be the Laver-Prikry forcing on A associated with the ideal $\text{fin}(A)$ of finite subsets of A . Let

$$D'_\alpha = \{T \cap A^{<\omega} : T \in D_\alpha \wedge T \cap A^{<\omega} \in \mathbb{L}_{\text{fin}(A)^*}\}.$$

Claim 1.5. $\{T_\alpha^n \cap A^{<\omega} : n \in \omega \text{ and } T_\alpha^n \cap A^{<\omega} \in \mathbb{L}_{\text{fin}(A)^*}\}$ is predense in $\mathbb{L}_{\text{fin}(A)^*}$. Therefore, D'_α is dense in $\mathbb{L}_{\text{fin}(A)^*}$.

Proof of Claim 1.5. Fix $\alpha < \kappa$. Let $S \in \mathbb{L}_{\text{fin}(A)^*}$ and $s = \text{stem}(S)$.

Then $\text{rk}_\alpha(s) < \infty$. Since $\{n \in \omega : s \frown \langle n \rangle \in T\} \in \text{fin}(A)^*$ and $H_s^\alpha \cap A$ is infinite, $H_s^\alpha \cap \{n \in \omega : t \frown \langle n \rangle \in T\} \neq \emptyset$.

By induction on rank, there exists $t \in S$ such that $t \supset s = \text{stem}(S)$ and $\text{rk}_\alpha(t) = 0$, that is, $t \in T_\alpha^n$ and $t \supset \text{stem}(T_\alpha^n)$ for some $n \in \omega$. Fix such $n \in \omega$.

Since $t \in A^{<\omega}$, $\text{stem}(T_\alpha^n) \in A^{<\omega}$. For every $u \in T_\alpha^n$ with $u \supset \text{stem}(T_\alpha^n)$, $A \subset^* \{n \in \omega : u \frown \langle n \rangle \in T_\alpha^n\}$. So $T_\alpha^n \cap A^{<\omega} \in \mathbb{L}_{\text{fin}(A)^*}$ and $T_\alpha^n \cap A^{<\omega}$ is compatible with S . Hence $\{T_\alpha^n \cap A^{<\omega} : n \in \omega \wedge T_\alpha^n \cap A^{<\omega} \in \mathbb{L}_{\text{fin}(A)^*}\}$ is predense. \square

Let $T_\alpha^{n'} = T_\alpha^n \cap A^{<\omega}$. For each $\alpha < \kappa$ and $n \in \omega$ with $T_\alpha^{n'} \in \mathbb{L}_{\text{fin}(A)^*}$, define $g_n^\alpha : A^{<\omega} \rightarrow \omega$ by

$$g_n^\alpha(s) = \begin{cases} \min\{n : A \setminus n \subset \text{Succ}_{T_\alpha^{n'}}(s)\} & \text{if } s \in T_\alpha^{n'} \text{ and } s \supset \text{stem}(T_\alpha^{n'}), \\ 0 & \text{otherwise.} \end{cases}$$

Notice that when $\text{Stem}(T_\alpha^n) \notin A^{<\omega}$, $T_\alpha^{n'} \notin \mathbb{L}_{\text{fin}(A)^*}$ and g_n^α is undefined.

Since $\kappa < \text{add}(\mathcal{M}) \leq \mathfrak{b}$, there exists $g : A^{<\omega} \rightarrow \omega$ such that for $\alpha < \kappa$ and $n \in \omega$, for almost all $t \in A^{<\omega}$, $g_n^\alpha(t) \leq g(t)$. Define $S \in \mathbb{L}_{\text{fin}(A)^*}$ so that

- (1) $\emptyset \in S$ and
- (2) if $s \in S$, then $s \frown \langle k \rangle \in S$ if and only if $k > g(s)$ and $k \in A$.

For each $\alpha < \kappa$, put

$$D''_\alpha = \{t \in S : \exists n \in \omega (t \in T''_\alpha, T''_\alpha \in \mathbb{L}_{\text{fin}(A)}^* \text{ and } \forall s \in S (s \supset t \rightarrow g(s) \geq g''_\alpha(s)))\}.$$

Let $M_\alpha = \{f \in [S] : \forall n \in \omega (f \upharpoonright n \notin D''_\alpha)\}$. Then M_α is nowhere dense in $[S]$. Since $\kappa < \text{add}(\mathcal{M}) \leq \text{cov}(\mathcal{M})$, there exists $f \in [S]$ such that $f \notin M_\alpha$ for every $\alpha < \kappa$.

Claim 1.6. *For every $\alpha < \kappa$, there exists $T \in D_\alpha$ such that $f \in [T]$.*

Proof. For $\alpha < \kappa$, let $n \in \omega$ such that $f \upharpoonright n \in D''_\alpha$. Then there exists $m \in \omega$ such that $f \upharpoonright n \in T''_\alpha$ and for $s \in S$ whenever $s \supset f \upharpoonright n$, $g(s) \geq g''_\alpha(s)$. By definition of g''_α , $S_{f \upharpoonright n} \subset T''_\alpha$. Hence $f \in [T''_\alpha]$. So $f \in [T]$ for some $T \in D_\alpha$. \square

By construction of f , f is a $\{D_\alpha : \alpha < \kappa\}$ -generic real, i.e., $\{T : f \in [T]\}$ is a filter intersecting with D_α for all $\alpha < \kappa$. \square

Corollary 1.7. *For every ideal \mathcal{I} on ω ,*

$$\mathfrak{m}(\mathbb{L}_{\mathcal{I}^*}) = \begin{cases} \min\{\mathfrak{b}, \text{sep}(\mathcal{I})\} & \text{if } \mathcal{I}^* \text{ is ultrafilter} \\ \min\{\text{add}(\mathcal{M}), \text{sep}(\mathcal{I})\} & \text{otherwise.} \end{cases}$$

1.2. Martin number of $\mathbb{M}_{\mathcal{I}^*}$. It seems that the rank argument does not work for the Mathias-Prikry type forcings. However, they can be investigated by studying the ideal $\mathcal{I}^{<\omega}$ on $[\omega]^{<\omega} \setminus \{\emptyset\}$ associated to an ideal \mathcal{I} on ω .

Definition 2. Given ideal \mathcal{I} on ω , let

$$\mathcal{I}^{<\omega} = \{A \subset [\omega]^{<\omega} \setminus \{\emptyset\} : \exists I \in \mathcal{I} \forall a \in A (a \cap I \neq \emptyset)\}.$$

This ideal was considered by Sirota [21] and Louveau [18] in the construction of an extremely disconnected topological group. Recall that an ultrafilter \mathcal{U} on ω is selective if for every partition $\{I_n : n \in \omega\}$ of ω either there is an n such that $I_n \in \mathcal{U}$ or there is a $U \in \mathcal{U}$ such that $|I_n \cap U| \leq 1$ for every $n \in \omega$.

Theorem 2.1. *For every ideal \mathcal{I} on ω ,*

$$\mathfrak{m}(\mathbb{M}_{\mathcal{I}^*}) = \begin{cases} \text{sep}(\mathcal{I}) & \text{if } \mathcal{I}^* \text{ is a selective ultrafilter.} \\ \min\{\text{sep}(\mathcal{I}^{<\omega}), \text{cov}(\mathcal{M})\} & \text{otherwise.} \end{cases}$$

If \mathcal{I}^* is a selective ultrafilter, then $\mathfrak{m}(\mathbb{M}_{\mathcal{I}^*}) = \text{cov}^*(\mathcal{I})$ by Theorem 1.2 (1).

To prove the rest of this theorem, we will first introduce two variations of $\text{sep}(\mathcal{I})$. Define $\widetilde{\text{sep}}(\mathcal{I})$ by $\kappa < \widetilde{\text{sep}}(\mathcal{I})$ if for $\mathcal{J} \subset \mathcal{I}$ and $\mathcal{H} \subset (\mathcal{I}^{<\omega})^+$ with $|\mathcal{J}| + |\mathcal{H}| \leq \kappa$, there exists $A \subset \omega$ such that

- (1) $|A \cap J| < \aleph_0$ for $J \in \mathcal{J}$ and
- (2) $|[A]^{<\omega} \cap H| = \aleph_0$ for $H \in \mathcal{H}$.

Define $\widetilde{\text{sep}}(\mathcal{I})$ by $\kappa < \widetilde{\text{sep}}(\mathcal{I})$ if for $\mathcal{J} \subset \mathcal{I}$ and $\mathcal{H} \subset (\mathcal{I}^{<\omega})^+$ with $|\mathcal{J}| + |\mathcal{H}| \leq \kappa$, there exists $A \subset \omega$ such that

- (1) $|A \cap J| < \aleph_0$ for $J \in \mathcal{J}$ and
- (2) $|[A \setminus n]^{<\omega} \cap H| = \aleph_0$ for $H \in \mathcal{H}$ and $n \in \omega$.

Claim 2.2. $\widetilde{\text{sep}}(\mathcal{I}) = \widetilde{\widetilde{\text{sep}}}(\mathcal{I})$.

Proof of Claim 2.2. By definition, it is clear that $\widetilde{\widetilde{\text{sep}}}(\mathcal{I}) \leq \widetilde{\text{sep}}(\mathcal{I})$. We shall show $\widetilde{\text{sep}}(\mathcal{I}) \geq \widetilde{\widetilde{\text{sep}}}(\mathcal{I})$.

Let $\mathcal{J} \subset \mathcal{I}$ and $\mathcal{H} \subset (\mathcal{I}^{<\omega})^+$ with $|\mathcal{J}| + |\mathcal{H}| < \widetilde{\text{sep}}(\mathcal{I})$. Let $\mathcal{H}^* = \{H_n : H \in \mathcal{H}, n \in \omega \text{ and } H_n = H \cap [\omega \setminus n]^{<\omega}\}$. Since $|\mathcal{J}| + |\mathcal{H}^*| = |\mathcal{J}| + |\mathcal{H}| < \widetilde{\text{sep}}(\mathcal{I})$, we can pick $A \subset \omega$ so that

- (1) $|A \cap J| < \aleph_0$ for $J \in \mathcal{J}$ and
- (2) $|[A]^{<\omega} \cap H| = \aleph_0$ for $H \in \mathcal{H}^*$.

Since $[A]^{<\omega} \cap H_n = [A]^{<\omega} \cap [\omega \setminus n]^{<\omega} \cap H = [A \setminus n]^{<\omega} \cap H$ for $H \in \mathcal{H}$ and $n \in \omega$, $|A \cap J| < \aleph_0$ for $J \in \mathcal{J}$ and $|[A \setminus n]^{<\omega} \cap H| = \aleph_0$ for $H \in \mathcal{H}$ and $n \in \omega$. Therefore $\widetilde{\text{sep}}(\mathcal{I}) \geq \widetilde{\widetilde{\text{sep}}}(\mathcal{I})$. \square

Lemma 2.3. *If \mathcal{I} is not a selective ultrafilter, then*

$$\mathfrak{m}(\mathbb{M}_{\mathcal{I}^*}) = \min\{\widetilde{\text{sep}}(\mathcal{I}), \text{cov}(\mathcal{M})\}.$$

Proof of Lemma 2.3. We shall show that $\mathfrak{m}(\mathbb{M}_{\mathcal{I}^*}) = \min\{\widetilde{\widetilde{\text{sep}}}(\mathcal{I}), \text{cov}(\mathcal{M})\}$.

(i) $\mathfrak{m}(\mathbb{M}_{\mathcal{I}^*}) \geq \min\{\widetilde{\widetilde{\text{sep}}}(\mathcal{I}), \text{cov}(\mathcal{M})\}$.

Let $\kappa < \widetilde{\widetilde{\text{sep}}}(\mathcal{I}), \text{cov}(\mathcal{M})$. Let $\{D_\alpha : \alpha < \kappa\}$ be a family of open dense subsets of $\mathbb{M}_{\mathcal{I}^*}$. Let $\{\langle s_\alpha^n, F_\alpha^n \rangle : n \in \omega\}$ be a maximal antichain in D_α . Let $I_\alpha^n = \omega \setminus F_\alpha^n \in \mathcal{I}$ for $n \in \omega$ and $\alpha < \kappa$. Let

$$H_s^\alpha = \{t \in [\omega]^{<\omega} : \exists n \in \omega (s_\alpha^n \subset s \cup t \subset s_\alpha^n \cup F_\alpha^n)\}.$$

Claim 2.4. $H_s^\alpha \in (\mathcal{I}^{<\omega})^+$ for all $s \in [\omega]^{<\omega}$ and $\alpha < \kappa$.

Proof of Claim 2.4. Let $s \in [\omega]^{<\omega}$, $\alpha < \kappa$ and $I \in \mathcal{I}$. Then $\langle s, \omega \setminus (I \cup s) \rangle \in \mathbb{M}_{\mathcal{I}^*}$. Since $\{\langle s_\alpha^n, F_\alpha^n \rangle : n \in \omega\}$ is a maximal antichain, $\langle s, \omega \setminus (I \cup s) \rangle$ is compatible with some $\langle s_\alpha^n, F_\alpha^n \rangle$. So there are $n \in \omega$, $t \in [\omega \setminus (I \cup s)]^{<\omega}$ and $F \in \mathcal{I}^*$ such that $\langle s \cup t, F \rangle \leq \langle s_\alpha^n, F_\alpha^n \rangle, \langle s, \omega \setminus (I \cup s) \rangle$.

Since $\langle s \cup t, F \rangle \leq \langle s_\alpha^n, F_\alpha^n \rangle$, $s_\alpha^n \subset s \cup t \subset s_\alpha^n \cup F_\alpha^n$. Since $\langle s \cup t, F \rangle \leq \langle s, \omega \setminus (I \cup s) \rangle$, $t \cap I = \emptyset$. Hence for each $I \in \mathcal{I}$, there exists $t \in H_s^\alpha$ such that $t \cap I = \emptyset$. Therefore $H_s^\alpha \in (\mathcal{I}^{<\omega})^+$. \square

As $\kappa < \widetilde{\widetilde{\text{sep}}}(\mathcal{I})$, there is an $A \subset \omega$ such that

- (1) $|A \cap I_\alpha^n| < \aleph_0$ for every $n \in \omega$ and $\alpha < \kappa$ and
- (2) $|[A \setminus n]^{<\omega} \cap H_s^\alpha| = \aleph_0$ for every $n \in \omega$, $s \in [\omega]^{<\omega}$ and $\alpha < \kappa$.

Let $\mathcal{A}_\alpha = \{\langle s_\alpha^n, F_\alpha^n \cap A \rangle : s_\alpha^n \subset A \text{ and } n \in \omega\} \subset \mathbb{M}_{\text{fin}(A)^*}$, where $\mathbb{M}_{\text{fin}(A)^*}$ is the Mathias-Prikry forcing associated with the ideal $\text{fin}(A)$ of finite subsets of A and $\mathbb{M}_{\text{fin}(A)^*}$ consists of pairs $\langle s, B \rangle$ such that $s \in [A]^{<\omega}$, $B \in \text{fin}(A)^*$ and $s \cap B = \emptyset$.

Claim 2.5. \mathcal{A}_α is predense in $\mathbb{M}_{\text{fin}(A)^*}$.

Proof of Claim 2.5. Let $\langle s, B \rangle \in \mathbb{M}_{\text{fin}(A)^*}$. Let $n \geq \max(s)$ such that $B \setminus n = A \setminus n$. Since $|[A \setminus n]^{<\omega} \cap H_s^\alpha| = \aleph_0$, pick $t \in [A \setminus n]^{<\omega} \cap H_s^\alpha$. Then there is an $n \in \omega$ such that $s_\alpha^n \subset s \cup t \subset s_\alpha^n \cup (F_\alpha^n \cap A)$. So $\langle s \cup t, (F_\alpha^n \setminus s \cup t) \cap A \rangle \leq \langle s_\alpha^n, F_\alpha^n \cap A \rangle$ and $\langle s \cup t, (F_\alpha^n \setminus s \cup t) \cap A \rangle \in \mathbb{M}_{\text{fin}(A)^*}$. Since $t \in [A \setminus n]^{<\omega} = [B \setminus n]^{<\omega}$, $\langle s \cup t, B \setminus (s \cup t) \rangle \leq \langle s, B \rangle$. So $\langle s, B \rangle$ is compatible with $\langle s_\alpha^n, F_\alpha^n \cap A \rangle$ for some $n \in \omega$. \square

Let $D'_\alpha = \{\langle s, F \cap A \rangle : s \subset A, F \cap A \in \text{fin}(A)^* \text{ and } \langle s, F \rangle \in D_\alpha\}$. Then D'_α is dense open subset of $\mathbb{M}_{\text{fin}(A)^*}$. Since $|\mathbb{M}_{\text{fin}(A)^*}| = \aleph_0$, $\mathbb{M}_{\text{fin}(A)^*} \cong \mathbb{C}$. Since $\kappa < \text{cov}(\mathcal{M})$, there exists A_{gen} such that for every $\alpha < \kappa$ there is $\langle s, F \cap A \rangle \in D'_\alpha$ so that $s \subset A_{\text{gen}} \subset s \cup (F \cap A)$. Hence, for every $\alpha < \kappa$ there exists $\langle s, F \rangle \in D_\alpha$ such that $s \subset A_{\text{gen}} \subset s \cup F$.

(ii) $\mathfrak{m}(\mathbb{M}_{\mathcal{I}^*}) \leq \min\{\widetilde{\text{sep}}(\mathcal{I}), \text{cov}(\mathcal{M})\}$.

Suppose $\kappa < \mathfrak{m}(\mathbb{M}_{\mathcal{I}^*})$. Let $\mathcal{J} \subset \mathcal{I}$ and $\mathcal{H} \subset (\mathcal{I}^{<\omega})^+$ with $|\mathcal{J}| + |\mathcal{H}| \leq \kappa$. Let $D_J = \{\langle s, F \rangle \in \mathbb{M}_{\mathcal{I}^*} : F \cap J = \emptyset\}$ for $J \in \mathcal{J}$, and let $E_H^n = \{\langle s, F \rangle \in \mathbb{M}_{\mathcal{I}^*} : |[s]^{<\omega} \cap H| \geq n\}$ for $H \in \mathcal{H}$ and $n \in \omega$. Since $\mathcal{J} \subset \mathcal{I}$ and $\mathcal{H} \subset (\mathcal{I}^{<\omega})^+$, D_J and E_H^n are dense subsets of $\mathbb{M}_{\mathcal{I}^*}$ for $J \in \mathcal{J}$, $H \in \mathcal{H}$ and $n \in \omega$. Let $A \subset \omega$ be a $\{D_J : J \in \mathcal{J}\} \cup \{E_H^n : H \in \mathcal{H} \wedge n \in \omega\}$ -generic real. Then $|A \cap J| < \aleph_0$ for $J \in \mathcal{J}$ and $|A \cap H| = \aleph_0$ for $H \in \mathcal{H}$. So $\kappa < \widetilde{\text{sep}}(\mathcal{I})$.

Since \mathcal{I}^* is not selective ultrafilter, $\mathbb{M}_{\mathcal{I}^*}$ adds a Cohen real (see [3]). Therefore $\kappa < \text{cov}(\mathcal{M})$. \square

Lemma 2.6.

$$\min\{\widetilde{\text{sep}}(\mathcal{I}), \text{cov}(\mathcal{M})\} = \min\{\text{sep}(\mathcal{I}^{<\omega}), \text{cov}(\mathcal{M})\}.$$

Proof. To prove $\min\{\widetilde{\text{sep}}(\mathcal{I}), \text{cov}(\mathcal{M})\} \geq \min\{\text{sep}(\mathcal{I}^{<\omega}), \text{cov}(\mathcal{M})\}$, we shall show that $\widetilde{\text{sep}}(\mathcal{I}) \geq \text{sep}(\mathcal{I}^{<\omega})$.

Claim 2.7. $\text{sep}(\mathcal{I}^{<\omega}) \leq \widetilde{\text{sep}}(\mathcal{I}) \leq \text{sep}(\mathcal{I})$.

Proof of Claim 2.7. Suppose $\kappa < \text{sep}(\mathcal{I}^{<\omega})$. Let $\mathcal{J} \subset \mathcal{I}$ and $\mathcal{H} \subset (\mathcal{I}^{<\omega})^+$ with $|\mathcal{J}| + |\mathcal{H}| \leq \kappa$. For $J \in \mathcal{J}$, put $\hat{J} = \{a \in [\omega]^{<\omega} : a \cap J \neq \emptyset\}$. Then $\hat{J} \in \mathcal{I}^{<\omega}$.

Let $A \subset [\omega]^{<\omega}$ be such that

- (1) $|A \cap \hat{J}| < \aleph_0$ for $J \in \mathcal{J}$ and
- (2) $|A \cap H| = \aleph_0$ for $H \in \mathcal{H}$.

Then $|\bigcup A \cap J| < \aleph_0$, and since $A \subset [\bigcup A]^{<\omega}$, $\bigcup A$ satisfies

- (1) $|\bigcup A \cap J| < \aleph_0$ for $J \in \mathcal{J}$ and
- (2) $|[\bigcup A]^{<\omega} \cap H| = \aleph_0$ for $H \in \mathcal{H}$.

Hence $\kappa < \widetilde{\text{sep}}(\mathcal{I})$. Therefore $\text{sep}(\mathcal{I}^{<\omega}) \leq \widetilde{\text{sep}}(\mathcal{I})$.

$\widetilde{\text{sep}}(\mathcal{I}) \leq \text{sep}(\mathcal{I})$ follows from the fact when $H \in \mathcal{I}^+$, $H^* = \{\{n\} : n \in H\} \in (\mathcal{I}^{<\omega})^+$. \square

To finish the proof of the theorem, we shall show $\min\{\widetilde{\text{sep}}(\mathcal{I}), \text{cov}(\mathcal{M})\} \leq \min\{\text{sep}(\mathcal{I}^{<\omega}), \text{cov}(\mathcal{M})\}$.

Suppose $\kappa < \widetilde{\text{sep}}(\mathcal{I})$, $\text{cov}(\mathcal{M})$. Let $\mathcal{J} \subset \mathcal{I}^{<\omega}$ and $\mathcal{H} \subset (\mathcal{I}^{<\omega})^+$ with $|\mathcal{J}| + |\mathcal{H}| \leq \kappa$. For $J \in \mathcal{J}$, fix $I_J \in \mathcal{I}$ so that $a \cap I_J \neq \emptyset$ for $a \in J$.

Let $A \subset \omega$ be such that

- (1) $|A \cap I_J| < \aleph_0$ for all $J \in \mathcal{J}$.
- (2) $|[A \setminus n]^{<\omega} \cap H| = \aleph_0$ for every $H \in \mathcal{H}$ and $n \in \omega$.

We will construct $B \subset [A]^{<\omega}$ so that

- (1) $|B \cap J| < \aleph_0$ for $J \in \mathcal{J}$.
- (2) $|B \cap H| = \aleph_0$ for $H \in \mathcal{H}$.

In order to do so, define a forcing notion \mathbb{P} by $\langle F, n \rangle \in \mathbb{P}$ if $F \in [[A]^{<\omega}]^{<\omega}$ and $n \in \omega$ ordered by $\langle F, n \rangle \leq \langle G, m \rangle$ if $F \supset G$, $n \geq m$ and $\min(a) \geq m$ for $a \in F \setminus G$. Since $|\mathbb{P}| = \aleph_0$, $\mathbb{C} \cong \mathbb{P}$. Let $D_{H,n}$ and $E_J \subset \mathbb{P}$ for $H \in \mathcal{H}$, $n \in \omega$ and $J \in \mathcal{J}$ be defined by

$$\begin{aligned} D_{H,n} &= \{\langle F, m \rangle : \exists a \in F (\min(a) > n \text{ and } a \in H)\}. \\ E_J &= \{\langle F, m \rangle : m > \max(A \cap I_J)\}. \end{aligned}$$

Then $D_{H,n}$ is dense for $H \in \mathcal{H}$ and $n \in \omega$, and E_J is dense for $J \in \mathcal{J}$.

Since $\kappa < \text{cov}(\mathcal{M})$, there is a $\{D_{H,n} : H \in \mathcal{H} \text{ and } n \in \omega\} \cup \{E_J : J \in \mathcal{J}\}$ -generic G . Let $A_G = \cup\{F : \langle F, n \rangle \in G\}$. Then

- (1) $|A_G \cap J| < \aleph_0$ for $J \in \mathcal{J}$ and
- (2) $|A_G \cap H| = \aleph_0$ for $H \in \mathcal{H}$.

So $\kappa < \text{sep}(\mathcal{I}^{<\omega})$. Therefore

$$\min\{\widetilde{\text{sep}}(\mathcal{I}), \text{cov}(\mathcal{M})\} \leq \min\{\text{sep}(\mathcal{I}^{<\omega}), \text{cov}(\mathcal{M})\}.$$

\square

Corollary 2.8. *For every ideal \mathcal{I} on ω ,*

$$\mathfrak{m}(\mathbb{M}_{\mathcal{I}^*}) = \begin{cases} \text{sep}(\mathcal{I}) & \text{if } \mathcal{I}^* \text{ is an ultrafilter.} \\ \min\{\text{sep}(\mathcal{I}^{<\omega}), \text{cov}(\mathcal{M})\} & \text{if } \mathcal{I} \text{ is not an ultrafilter.} \end{cases}$$

3. PRESERVATION PROPERTIES OF $\mathbb{M}_{\mathcal{I}^*}$

The methods for studying properties of the forcing $\mathbb{L}_{\mathcal{I}^*}$ are well known (see [2, 7, 12]). Here we concentrate on the preservation properties of the forcings $\mathbb{M}_{\mathcal{I}^*}$. In [12] it is shown that a useful characterization for when $\mathbb{L}_{\mathcal{I}^*}$ preserves ω -hitting families. An analogous result also holds for $\mathbb{M}_{\mathcal{I}^*}$.¹

Theorem 3.1. *Let \mathcal{I} be an ideal on ω . The following are equivalent:*

- (1) $\forall X \in (\mathcal{I}^{<\omega})^+ \forall \mathcal{J} \leq_K \mathcal{I}^{<\omega} \upharpoonright X$ (\mathcal{J} is not ω -hitting,)
- (2) $\mathbb{M}_{\mathcal{I}^*}$ strongly preserves ω -hitting families, and
- (3) $\mathbb{M}_{\mathcal{I}^*}$ preserves ω -hitting families.

Proof. From (1) to (2).

Suppose (2) doesn't hold. Let \dot{A} be $\mathbb{M}_{\mathcal{I}^*}$ -names witnessing the negation of (2), i.e., for every $(B_n : n \in \omega)$ there exists $B \in [\omega]^\omega$ such that $|B_n \cap B| = \aleph_0$ for every $n \in \omega$ and there exist $p_B = \langle s_B, F_B \rangle \in \mathbb{M}_{\mathcal{I}^*}$ and $m_B \in \omega$ such that $p_B \Vdash B \cap \dot{A} \subset m_B$.

Let \mathcal{B} be the family of all such B . By the assumption that \mathcal{B} is ω -hitting, there are $s \in [\omega]^{<\omega}$ and $m \in \omega$ such that $\mathcal{B}_0 = \{B \in \mathcal{B} : s_B = s \text{ and } m_B = m\}$ is ω -hitting (If an ω -hitting family is split into countably pieces, one of them is ω -hitting). Fix such $s \in [\omega]^{<\omega}$ and $m \in \omega$ and let

$$X_s = \{t \in [\omega]^{<\omega} : \exists k > m \exists F \in \mathcal{I}^* (\langle s \cup t, F \rangle \Vdash k \in \dot{A})\}.$$

Claim 3.2. $X_s \in (\mathcal{I}^{<\omega})^+$.

Proof of Claim 3.2. Given $I \in \mathcal{I}$, there are $t \in [\omega \setminus I]^{<\omega}$, $k > m$ and $F \in \mathcal{I}$ such that $\langle s \cup t, F \rangle \leq \langle s, \omega \setminus I \rangle$ and $\langle s \cup t, F \rangle \Vdash k \in \dot{A}$. Then $t \in X_s$ and $t \cap I = \emptyset$. \square

Define $f : X_s \rightarrow \omega$ by

$$f(t) = \begin{cases} \max\{k > m : \exists F \in \mathcal{I}^* (\langle s \cup t, F \rangle \Vdash k \in \dot{A})\} \\ \quad \text{if there are finitely many such } k, \\ \min\{k > \max(t \cup \{m\}) : \exists F \in \mathcal{I}^* (\langle s \cup t, F \rangle \Vdash k \in \dot{A})\} \\ \quad \text{otherwise.} \end{cases}$$

Claim 3.3. *For every $k \in \omega$, $f^{-1}[\omega \setminus k] \in (\mathcal{I}^{<\omega})^+$.*

¹A forcing \mathbb{P} strongly preserves ω -hitting families if given a \mathbb{P} -name \dot{A} for an infinite subset of ω there is a countable family \mathcal{H} of infinite subsets of ω such that whenever $B \subseteq \omega$ has an infinite intersection with every element of \mathcal{H} then $\Vdash_{\mathbb{P}} “|\dot{A} \cap B| = \omega”$.

So we can assume that for all but finitely many $k > m$, $f^{-1}(\{k\}) \in \mathcal{I}^{<\omega}$ or there exist infinitely many $k \in \omega$ such that $f^{-1}(\{k\}) \in (\mathcal{I}^{<\omega})^+$.

Claim 3.4. *For all but finitely many $k > m$, $f^{-1}(\{k\}) \in \mathcal{I}^{<\omega}$.*

Proof. Assume to the contrary that there are infinitely many $k > m$ such that $f^{-1}(\{k\}) \in (\mathcal{I}^{<\omega})^+$. Let $C = \{k > m : f^{-1}(\{k\}) \in (\mathcal{I}^{<\omega})^+\}$. Since \mathcal{B}_0 is ω -hitting, there exists $B \in \mathcal{B}_0$ such that $B \cap C$ is infinite. Let $k > m$ such that $k \in C \cap B$ and $f^{-1}(\{k\}) \in (\mathcal{I}^{<\omega})^+$. Then $|[F_B]^{<\omega} \cap f^{-1}(\{k\})| = \aleph_0$. Let $t \in [F_B]^{<\omega} \cap f^{-1}(\{k\})$. Then there exists $F \in \mathcal{I}^*$ such that $\langle s \cup t, F \rangle \Vdash k \in \dot{A} \cap B$. Since $t \in [F_B]^{<\omega}$, $\langle s, F_B \rangle$ is compatible with $\langle s \cup t, F \rangle$. However $\langle s, F_B \rangle \Vdash \dot{A} \cap B \subset m$, which is a contradiction. \square

Claim 3.5. *f witnesses that $\langle \mathcal{B}_0 \rangle \leq_K \mathcal{I}^{<\omega} \upharpoonright X$.*

Proof. Assume to the contrary that there is a $B \in \mathcal{B}_0$ such that $f^{-1}[B] \in (\mathcal{I}^{<\omega})^+$. Since $F_B \in \mathcal{I}^*$ and $|[F_B]^{<\omega} \cap f^{-1}[B]| = \aleph_0$, there is a $t \in [F_B]^{<\omega} \cap f^{-1}[B]$ such that for some $F \in \mathcal{I}^*$ and $k > m$ $\langle s \cup t, F \rangle \Vdash k \in \dot{A} \cap B$.

Since $t \in [F_B]^{<\omega}$, $s \subset s \cup t \subset s \cup F_B$. So $\langle s, F_B \rangle$ is compatible with $\langle s \cup t, F \rangle$. However, $\langle s, F_B \rangle \Vdash \dot{A} \cap B \subset m$, which is a contradiction. \square

Since $\langle \mathcal{B}_0 \rangle$ is ω -hitting, (1) doesn't hold.

Obviously (2) implies (3).

We shall prove (3) implies (1). Assume to the contrary that there exists an ideal \mathcal{J} on ω such that there exist $X \in (\mathcal{I}^{<\omega})^+$ and $f : X \rightarrow \omega$ so that

- (1) For every $J \in \mathcal{J}$, $f^{-1}[J] \in \mathcal{I}^{<\omega}$ and
- (2) \mathcal{J} is ω -hitting.

Let \dot{a}_{gen} be the canonical name for the $\mathbb{M}_{\mathcal{I}^*}$ -generic subset of ω . We shall show that $\Vdash \mathcal{J}$ is not ω -hitting. Let \dot{X}_n be a $\mathbb{M}_{\mathcal{I}^*}$ -name such that

$$\Vdash \dot{X}_n = f[[\dot{a}_{gen} \setminus n]^{<\omega}].$$

Claim 3.6. $\Vdash \dot{X}_n$ is infinite.

Proof of the Claim. If $\Vdash \dot{X}_n$ were finite, then

$$\Vdash [\dot{a}_{gen} \setminus n]^{<\omega} \cap X \subset f^{-1}[\dot{X}_n] \in \mathcal{I}^{<\omega}.$$

However, $\Vdash \forall I \in \mathcal{I} (|\dot{a}_{gen} \cap I| < \aleph_0)$ and $\Vdash |[\dot{a}_{gen} \setminus n]^{<\omega} \cap X| = \aleph_0$ by genericity. So $\Vdash \forall I \in \mathcal{I} \forall n \in \omega \exists a \in [\dot{a}_{gen} \setminus n]^{<\omega} (a \cap I = \emptyset)$, which is a contradiction. \square

Claim 3.7. $\Vdash \forall J \in \mathcal{J} \exists n \in \omega (J \cap \dot{X}_n = \emptyset)$.

Proof. For every $J \in \mathcal{J}$ and $\langle s, F \rangle \in \mathbb{M}_{\mathcal{I}^*}$, pick $I \in \mathcal{I}$ such that $a \cap I \neq \emptyset$ for $a \in f^{-1}[J]$, $G = F \cap (\omega \setminus I)$ and $n = \max(s)$. Then $\langle s, G \rangle \Vdash [\dot{a}_{gen} \setminus n]^{<\omega} \cap f^{-1}[J] = \emptyset$. So $\langle s, G \rangle \Vdash \dot{X}_n \cap J = \emptyset$. \square

So \mathcal{J} is not ω -hitting in the extension, contradiction. \square

Now, we turn our attention to the question of when does the forcing $\mathbb{M}_{\mathcal{I}^*}$ add a dominating real. This line of investigation was started by M. Canjar in [14], where he assuming $\mathfrak{d} = \mathfrak{c}$ constructed an ultrafilter \mathcal{U} such that $\mathbb{M}_{\mathcal{U}}$ doesn't add any dominating reals. He also noticed that such an ultrafilter has to be a P-point without rapid Rudin-Keisler predecessors (see e.g. [1] for definitions and more information) and asked whether the converse is also true. Here we give a simple combinatorial characterizations of ideals \mathcal{I} such that $\mathbb{M}_{\mathcal{I}^*}$ doesn't add any dominating reals.

We call an ideal \mathcal{J} a P^+ -ideal if for every decreasing sequence $\{X_n : n \in \omega\}$ of \mathcal{J} positive sets, there is an $X \in \mathcal{J}^+$ such that $X \subset^* X_n$ for all $n \in \omega$.

Theorem 3.8. *The following are equivalent.*

- (1) $\mathbb{M}_{\mathcal{I}^*}$ adds a dominating real.
- (2) $\mathcal{I}^{<\omega}$ is not a P^+ -ideal.

Proof. (1) implies (2).

Let \dot{g} be a $\mathbb{M}_{\mathcal{I}^*}$ -name for a dominating real, i.e., $\forall f \in \omega^\omega \cap V (\Vdash f <^* \dot{g})$. In particular, for every $f \in \omega^\omega \cap V$, there are $s_f \in [\omega]^{<\omega}$, $F_f \in \mathcal{I}^*$ and $n_f \in \omega$ such that

$$\langle s_f, F_f \rangle \Vdash \forall n \geq n_f (f(n) < \dot{g}(n)).$$

If one partitions a dominating family into countably many pieces, one of the pieces is also dominating. So there are $s \in [\omega]^{<\omega}$ and $n \in \omega$ such that

$$\mathcal{F} = \{f \in \omega^\omega : s_f = s \wedge n_f = n\}$$

is a dominating family. Fix such $s \in [\omega]^{<\omega}$ and $n \in \omega$ and let

$$X_s = \{t \in [\omega \setminus \max(s)]^{<\omega} : \exists F \in \mathcal{I}^* \exists m \geq n (\langle s \cup t, F \rangle \text{ decides } \dot{g}(m))\}.$$

Claim 3.9. $X_s \in (\mathcal{I}^{<\omega})^+$.

For each $t \in X_s$, let $z_t = \{m \geq n : \exists F \in \mathcal{I}^* (\langle s \cup t, F \rangle \text{ decides } \dot{g}(m))\}$. Put $k_t, l_t \in \omega$ so that

$$k_t = \begin{cases} \max(z_t) & \text{if } |z_t| < \omega \\ \min(z_t \setminus \max(t)) & \text{otherwise.} \end{cases}$$

and there is an $F \in \mathcal{I}^*$ such that $\langle s \cup t, F \rangle \Vdash \dot{g}(k_t) = l_t$.

Then define $H : X_s \rightarrow \omega \times \omega$ by $H(t) = \langle k_t, l_t \rangle$.

Claim 3.10. For every $m \in \omega$, $H^{-1}[(\omega \setminus m) \times \omega] \in (\mathcal{I}^{<\omega})^+$.

Let $K = \{k_t : t \in X_s\}$. Then K is infinite and let $\{k_i : i \in \omega\}$ be its increasing enumeration. Put $K_i = \{l \in \omega : \langle k_i, l \rangle \in H[X_s]\}$.

Claim 3.11. There are infinitely many $i \in \omega$ such that K_i is infinite.

Proof of Claim 3.11. Assume to the contrary that for all but finitely many $i \in \omega$, K_i is finite. Then define $g : \omega \rightarrow \omega$ by

$$g(m) = \begin{cases} \max(K_i) & \text{if } m = k_i \text{ and } |K_i| < \aleph_0 \text{ for some } i \in \omega, \\ 0 & \text{otherwise.} \end{cases}$$

Since \mathcal{F} is a dominating family, there is an $f \in \mathcal{F}$ such that $g \leq^* f$. Let $m_0 \geq n$ be such that $g(m) \leq f(m)$ for $m \geq m_0$ and $k_i \geq m_0$ implies $|K_i| < \aleph_0$.

By Claim 3.10, $H^{-1}[(\omega \setminus m) \times \omega] \cap [F]^{<\omega} \in (\mathcal{I}^{<\omega})^+$ for $F \in \mathcal{I}^*$ and $m \in \omega$.

Let $t \in [F_f]^{<\omega} \cap H^{-1}[(\omega \setminus m_0) \times \omega]$. Then there is an $F \in \mathcal{I}^*$ such that $\langle s \cup t, F \rangle \Vdash \dot{g}(k_t) = l_t \leq g(k_t) \leq f(k_t)$. However, $\langle s, F_f \rangle$ is compatible with $\langle s \cup t, F \rangle$ and $\langle s, F_f \rangle \Vdash f(k_t) < \dot{g}(k_t)$, which is a contradiction. \square

Without loss of generality, we can assume that for every $i \in \omega$, K_i is infinite.

Let $Y_m = H^{-1}[\bigcup_{i>m} K_i]$ for $m \in \omega$. Then $Y_{m+1} \subset Y_m$. As Claim 3.10, we can prove the following.

Claim 3.12. $Y_m \in (\mathcal{I}^{<\omega})^+$ for $m \in \omega$.

Proof of Claim 3.12. Let $I \in \mathcal{I}$ and $m \in \omega$. We shall show that there exists $t \in Y_m$ such that $t \cap I = \emptyset$. Let $t \in [\omega \setminus I]^{<\omega}$ such that $\langle s \cup t, F \rangle \leq \langle s, \omega \setminus I \rangle$ decides $\dot{g}(k)$ for some $k > m$ and $\max(t) > m$. By definition of k_t , $k_t \geq k > m$. Then $H(t) \in \bigcup_{i>m} K_i$. So $t \in Y_m$ and $t \cap I = \emptyset$. \square

Let $Y \subset^* Y_m$ for $m \geq n$. We shall show that $Y \in \mathcal{I}^{<\omega}$.

Assume to the contrary that $Y \in (\mathcal{I}^{<\omega})^+$.

Since $Y \subset^* Y_m$, $K_m \cap H[Y]$ is finite for every $m \in \omega$. Define a function h from ω to ω by

$$h(m) = \begin{cases} \max\{l_t : \exists t \in Y \cap Y_m\} & \text{if } Y \cap Y_m \neq \emptyset, \\ 0 & \text{otherwise.} \end{cases}$$

Then there are infinitely many m such that $h(m) > 0$.

Since \mathcal{F} is a dominating family, there is an $f \in \mathcal{F}$ such that $h \leq^* f$. Let $m_0 \geq n$ be such that $h(m) \leq f(m)$ for $m \geq m_0$. Since $Y \subset^* Y_m$, $F_f \in \mathcal{I}^*$ and $Y \in (\mathcal{I}^{<\omega})^+$, there is an $m \geq m_0$ such that $Y \cap Y_m \cap$

$[F_f]^{<\omega} \neq \emptyset$. Let $t \in Y \cap Y_m \cap [F_f]^{<\omega}$. Since $t \in Y$ there is an $F \in \mathcal{I}^*$ such that $\langle s \cup t, F \rangle \Vdash \dot{g}(m) \leq h(m)$. However, $\langle s, F_f \rangle \Vdash \text{“}\forall m \geq n(f(m) < \dot{g}(m))\text{”}$ and $\langle s \cup t, F \rangle$ is compatible with $\langle s, F_f \rangle$, which is a contradiction. Therefore $Y \in \mathcal{I}^{<\omega}$. So $\mathcal{I}^{<\omega}$ is not P^+ -ideal.

(2) implies (1).

Let $\langle X_n : n \in \omega \rangle$ be a decreasing sequence of $\mathcal{I}^{<\omega}$ -positive sets without a pseudointersection in $(\mathcal{I}^{<\omega})^+$. Let $\langle a_k : k \in \omega \rangle$ be an enumeration of $[\omega]^{<\omega} \setminus \{\emptyset\}$ and let \dot{a}_{gen} be the canonical name for the $\mathbb{M}_{\mathcal{I}^*}$ -generic real. Define a $\mathbb{M}_{\mathcal{I}^*}$ -name \dot{g} for a function from ω to ω by

$$\begin{aligned} \Vdash \dot{g}(n) = \min\{k : a_k \subset [\dot{a}_{gen}]^{<\omega} \cap X_n \wedge \\ \max(\bigcup\{a_m : l < n \wedge m = \dot{g}(l)\}) < \min(a_k)\}. \end{aligned}$$

We shall show that \dot{g} is a dominating real. Let $f \in \omega^\omega \cap V$ and $\langle s, F \rangle \in \mathbb{M}_{\mathcal{I}^*}$. Let $I_f = \{a_k \in [\omega]^{<\omega} \setminus \{\emptyset\} : \exists n \in \omega (a_k \in X_n \wedge k \leq f(n))\}$. Then $I_f \subset^* X_n$ for every $n \in \omega$. Therefore $I_f \in \mathcal{I}^{<\omega}$ by definition of X_n . Let $I \in \mathcal{I}$ such that $\forall a \in I_f (a \cap I \neq \emptyset)$. Then $F \setminus I \in \mathcal{I}^*$ and $[F \setminus I]^{<\omega} \cap I_f = \emptyset$.

Claim 3.13. *Let $\langle t_n : n < \alpha \rangle$ be a sequence of finite subsets of ω such that*

- (1) $t_n \in [s \cup (F \setminus I)]^{<\omega} \cap X_n$
- (2) $\max(t_n) < \min(t_{n+1})$
- (3) $\exists k \in \omega (t_n = a_k \wedge k \leq f(n))$

Then $\alpha \leq |s|$.

Proof of Claim. If $t \in [F \setminus I]^{<\omega}$, then $t = a_k$ and $t \in X_n$ implies $k > f(n)$ by $[F \setminus I]^{<\omega} \cap I_f = \emptyset$. So by (2), $\alpha \leq |s|$. \square

Put $|s|=m$. Then $\langle s, F \setminus I \rangle \leq \langle s, F \rangle$ and

$$\langle s, F \setminus I \rangle \Vdash \forall n > m (f(n) < \dot{g}(n)).$$

\square

Recently, using our characterizations, M. Hrušák and J. Verner showed that if \mathcal{I} is an F_σ P -ideal, then $\mathcal{P}(\omega)/\mathcal{I}$ adds an ultrafilter \mathcal{U} which is a P -point without rapid RK -predecessors, but \mathcal{U}^* is not a P^+ -ideal [17]. Thus this answers Canjar's question in the negative. Moreover, A. Blass, M. Hrušák and J. Verner [4] (also using our theorem) showed that $\mathbb{M}_{\mathcal{U}}$ doesn't add any dominating reals if and only if \mathcal{U} is strong P -point.

4. CONCLUDING REMARKS AND OPEN PROBLEMS.

It is still interesting to try to better understand ideals \mathcal{I} for which $\mathbb{M}_{\mathcal{I}^*}$ doesn't add any dominating reals. An interesting class of ideals in this respect are those generated by maximal almost disjoint (mad) families.

Theorem 4.1. [6] *Assume $\mathfrak{b} = \mathfrak{c}$. Then there exists a mad family \mathcal{A} such that $\mathbb{M}_{\mathcal{I}(\mathcal{A})}$ adds a dominating real.*

Question 4.2. [6] *Is it consistent that there is no mad family \mathcal{A} such that $\mathbb{M}_{\mathcal{I}(\mathcal{A})}$ adds a dominating real?*

As far as definable ideals are concern, J. Brendle has in [6] that an $\mathbb{M}_{\mathcal{I}^*}$ doesn't add any dominating reals for any F_σ -ideal \mathcal{I} . This follows directly from our characterization. However, it is not clear whether this characterizes F_σ -ideals among Borel ones:

Question 4.3. *Is there a Borel ideal \mathcal{I} which is not F_σ , yet $\mathbb{M}_{\mathcal{I}^*}$ doesn't add any dominating reals?*

However, we have the following useful approximation:

Theorem 4.4. *Suppose \mathcal{I} is a Borel ideal. Then the following are equivalent.*

- (1) \mathcal{I} can be extended to an ideal \mathcal{J} such that $\mathbb{M}_{\mathcal{J}^*}$ which doesn't add any dominating reals.
- (2) \mathcal{I} can be extended to a P^+ -ideal.
- (3) \mathcal{I} can be extended to an F_σ -ideal.

Proof. (3) implies (1).

This is proved in Brendle's paper [6], but it also follows from our theorem using the following two simple observations:

- (i) If \mathcal{I} is F_σ then $\mathcal{I}^{<\omega}$ is F_σ , and
- (ii) every F_σ -ideal is P^+ .

(1) implies (2).

Suppose (2) doesn't hold. Then every \mathcal{J} extending \mathcal{I} is not P^+ .

Claim 4.5. *If $\mathcal{J}^{<\omega}$ is P^+ , then \mathcal{J} is P^+ .*

Proof of Claim. Let $\{Y_n : n \in \omega\}$ be a decreasing sequence of \mathcal{J}^+ . Put $Y_n^* = \{\{k\} : k \in Y_n\}$ for $n < \omega$. Then $Y_n^* \in (\mathcal{J}^{<\omega})^+$. By assumption, there exists $Y^* \in (\mathcal{J}^{<\omega})^+$ such that $Y^* \subset^* Y_n^*$ for $n < \omega$. Put $Y = \bigcup Y^*$. Then $Y \in \mathcal{J}^+$ and $Y \subset^* Y_n$ for $n < \omega$. So \mathcal{J} is P^+ . \square

By this claim, $\mathcal{J}^{<\omega}$ is not P^+ . So $\mathbb{M}_{\mathcal{J}^*}$ adds a dominating real by our theorem.

(2) implies (3).

This follows from a theorem of D. Meza and M. Hrušák (see [16]). \square

This, in particular, shows that the ideal \mathcal{Z} of sets Banach density zero can not be extended to an ideal \mathcal{I} such that $\mathbb{M}_{\mathcal{I}^*}$ doesn't add any dominating reals, as it can not be extended to an F_σ -ideal.

Corollary 4.6. *Let \mathcal{Z} be the density zero ideal. If $\mathcal{Z}^* \subset \mathcal{F}$, then $\mathbb{M}_{\mathcal{F}}$ adds a dominating real.*

Question 4.7. *Is there forcing notion \mathbb{P} which destroys \mathcal{Z} and doesn't add a dominating real?*

ACKNOWLEDGMENT

We thank David Meza-Alcántara, Jörg Brendle, Hiroshi Sakai and Lyubomyr Zdomskyy for stimulating discussions.

REFERENCES

- [1] Tomek Bartoszyński and Haim Judah. *Set theory. On the structure of the real line*. A K Peters, Wellesley, MA, 1995.
- [2] James E. Baumgartner and Peter Dordal. Adjoining dominating functions. *Journal of Symbolic Logic*, 50(1):94–101, 1985.
- [3] Aleksander Błaszczyk and Saharon Shelah. Regular subalgebras of complete boolean algebras. *Journal of Symbolic Logic*, 66(2):792–800, 2001.
- [4] Andreas Blass, Michael Hrušák, and Jonathan Verner. On strong p -points. preprint.
- [5] Andreas Blass and Saharon Shelah. Ultrafilters with small generating sets. *Israel Journal of Mathematics*, 65(3):259–271, 1989.
- [6] Jörg Brendle. Mob families and mad families. *Archive for Mathematical Logic*, 37(3):183–197, 1997.
- [7] Jörg Brendle. Van douwen's diagram for dense sets of rationals. *Annals of Pure and Applied Logic*, 143(1-3):54–69, 2006.
- [8] Jörg Brendle. *Algebra, Logic and Set Theory*, volume 4 of *Studies in Logic*, chapter Independence for distributivity numbers, pages 63–84. College Publications, London, 2007.
- [9] Jörg Brendle. Distinguishing groupwise density numbers. *Monatshefte für Mathematik*, 152(3):207–215, 2007.
- [10] Jörg Brendle. Mad families and ultrafilters. *Acta Universitatis Carolinae. Mathematica et Physica*, 48(2):19–35, 2007.
- [11] Jörg Brendle and Vera Fisher. Mad families, splitting families, and large continuum. *The Journal of Symbolic Logic*, 76(1):198–208, 2011.
- [12] Jörg Brendle and Michael Hrušák. Countable Fréchet boolean groups: An independence result. *The Journal of Symbolic Logic*, 74(3):1061–1068, 2009.

- [13] Jörg Brendle and Saharon Shelah. Ultrafilters on ω —their ideals and their cardinal characteristics. *Transactions of the American Mathematical Society*, 351(1-2):143–161, 1999.
- [14] R. Michael Canjar. Mathias forcing which does not add dominating reals. *Proceedings of the American Mathematical Society*, 104(4):1239–1248, 1988.
- [15] Fernando Hernández-Hernández and Michael Hrušák. Cardinal invariants of analytic P -ideals. *Canadian Journal of Mathematics*, 59(3):575–595, 2007.
- [16] Michael Hrušák. *Set theory and its applications*, volume 533 of *Contemporary Mathematics*, chapter Combinatorics of filters and ideals, pages 29–69. Amer Mathematical Society, 2011.
- [17] Michael Hrušák and Jonathan L. Verner. Adding ultrafilters by definable quotients. preprint.
- [18] Alain Louveau. Sur un article de S. Sirota. *Bulletin des Sciences Mathématiques. 2e Série*, 96:3–7, 1972.
- [19] Dilip Raghavan. A model with no strongly separable almost disjoint families. *Israel Journal of Mathematics*, to appear.
- [20] Saharon Shelah. Two cardinal invariants of the continuum ($\mathfrak{d} < \mathfrak{a}$) and fs linearly ordered iterated forcing. *Acta Mathematica*, 192(2):187–223, 2004.
- [21] S. M. Sirota. A product of topological groups, and extremal disconnectedness. *Matematicheskij Sbornik. Novaya Seriya*, 79(121):179–192, 1969.

INSTITUTO DE MATEMÁTICAS, UNAM, APARTADO POSTAL 61-3, XANGARI,
58089, MORELIA, MICHOACÁN, MÉXICO.

E-mail address: michael@matmor.unam.mx

E-mail address: minami@kurt.scitec.kobe-u.ac.jp